


Child Welfare Trauma Training

for Behavioral Health Providers



1



Child Welfare Trauma Training:

Creating Trauma-Informed Child Welfare Practice: Introduction to the Essential Elements



Trauma-Informed Child Welfare Practice

The trauma-informed provider:

- Understands the impact of trauma on a child's behavior, development, relationships, and survival strategies
- Can integrate that understanding into planning for the child and family
- Understands his or her role in responding to child traumatic stress

Trauma-Informed Child Welfare Practice, cont'd

The Essential Elements:

- Are the province of ALL professionals who work in and with the child welfare system
- Must, when implemented, take into consideration the child's developmental level and reflect sensitivity to the child's family, culture, and language
- Help child welfare systems achieve the goals of safety, permanency and well-being

Essential Elements of Trauma-Informed Child Welfare Practice


1. Maximize the child's sense of safety.
2. Assist children in reducing overwhelming emotion.
3. Help children make new meaning of their trauma history and current experiences.
4. Address the impact of trauma and subsequent changes in the child's behavior, development, and relationships.
5. Coordinate services with other agencies.

Essential Elements of Trauma-Informed Child Welfare Practice

6. Utilize comprehensive assessment of the child's trauma experiences and their impact on the child's development and behavior to guide services.
7. Support and promote positive and stable relationships in the life of the child.
8. Provide support and guidance to child's family and caregivers.
9. Manage professional and personal stress.

Child Welfare Trauma Training :

What Is Child Traumatic Stress?



Artwork courtesy of the International Child Art Foundation (www.icaf.org)

What Is Child Traumatic Stress?

- Child traumatic stress refers to the *physical and emotional responses* of a child to events that threaten the life or physical integrity of the child or of someone critically important to the child (such as a parent or sibling).
- Traumatic events overwhelm a child's capacity to cope and elicit feelings of terror, powerlessness, and out-of-control physiological arousal.

What Is Child Traumatic Stress, cont'd

- A child's response to a traumatic event may have a profound effect on his or her perception of self, the world, and the future.
- Traumatic events may affect a child's:
 - Ability to trust others
 - Sense of personal safety
 - Effectiveness in navigating life changes

Types of Traumatic Stress

- **Acute trauma** is a single traumatic event that is limited in time. Examples include:
 - Serious accidents
 - Community violence
 - Natural disasters (earthquakes, wildfires, floods)
 - Sudden or violent loss of a loved one
 - Physical or sexual assault (e.g., being shot or raped)
- During an acute event, children go through a variety of feelings, thoughts, and physical reactions that are frightening in and of themselves and contribute to a sense of being overwhelmed.

10

Types of Traumatic Stress, cont'd

- **Chronic trauma** refers to the experience of multiple traumatic events.
- These may be multiple and varied events—such as a child who is exposed to domestic violence, is involved in a serious car accident, and then becomes a victim of community violence—or longstanding trauma such as physical abuse, neglect, or war.
- The effects of chronic trauma are often cumulative, as each event serves to remind the child of prior trauma and reinforce its negative impact.

Types of Traumatic Stress, cont'd

- **Complex trauma** describes both exposure to chronic trauma—usually caused by adults entrusted with the child’s care—and the impact of such exposure on the child.
- Children who experienced complex trauma have endured multiple interpersonal traumatic events from a very young age.
- Complex trauma has profound effects on nearly every aspect of a child’s development and functioning.

Source: Cook et al. (2005). *Psychiatr Ann*, 35(5):390-398.



Prevalence of Trauma—United States

- One in four children/adolescents experience at least one potentially traumatic event before the age of 16.¹
- In a 1995 study, 41% of middle school students in urban school systems reported witnessing a stabbing or shooting in the previous year.²
- Four out of 10 U.S. children report witnessing violence; 8% report a lifetime prevalence of sexual assault, and 17% report having been physically assaulted.³

1. Costello et al. (2002). *J Traum Stress*;5(2):99-112.

2. Schwab-Stone et al. (1995). *J Am Acad Child Adolesc Psychiatry*;34(10):1343-1352.

3. Kilpatrick et al. (2003). US Dept. Of Justice. <http://www.ncjrs.gov/pdffiles1/nij/194972.pdf>.

Prevalence of Trauma in the Child Welfare Population

- A national study of adult “foster care alumni” found higher rates of PTSD (21%) compared with the general population (4.5%). This was higher than rates of PTSD in American war veterans.¹
- Nearly 80% of abused children face at least one mental health challenge by age 21.²

1. Pecora, et al. (December 10, 2003). *Early Results from the Casey National Alumni Study*. Available at: http://www.casey.org/NR/rdonlyres/CEFBB1B6-7ED1-440D-925A-E5BAF602294D/302/casey_alumni_studies_report.pdf.

2. ASTHO. (April 2005). *Child Maltreatment, Abuse, and Neglect*. Available at: <http://www.astho.org/pubs/Childmaltreatmentfactsheet4-05.pdf>.

Other Sources of Ongoing Stress

- Children in the child welfare system frequently face other sources of ongoing stress that can challenge workers' ability to intervene. Some of these sources of stress include:
 - Poverty
 - Discrimination
 - Separations from parent/siblings
 - Frequent moves
 - School problems
 - Traumatic grief and loss
 - Refugee or immigrant experiences

Effects of Trauma Exposure on Children

- When trauma is associated with the failure of those who should be protecting and nurturing the child, it has profound and far-reaching effects on nearly every aspect of the child's life.
- Children who have experienced the types of trauma that precipitate entry into the child welfare system typically suffer impairments in many areas of development and functioning, including:

Effects of Trauma Exposure, cont'd

- **Attachment.** Traumatized children feel that the world is uncertain and unpredictable. They can become socially isolated and can have difficulty relating to and empathizing with others.
- **Biology.** Traumatized children may experience problems with movement and sensation, including hypersensitivity to physical contact and insensitivity to pain. They may exhibit unexplained physical symptoms and increased medical problems.
- **Mood regulation.** Children exposed to trauma can have difficulty regulating their emotions as well as difficulty knowing and describing their feelings and internal states.

Effects of Trauma Exposure, cont'd

- **Dissociation.** Some traumatized children experience a feeling of detachment or depersonalization, as if they are “observing” something happening to them that is unreal.
- **Behavioral control.** Traumatized children can show poor impulse control, self-destructive behavior, and aggression towards others.
- **Cognition.** Traumatized children can have problems focusing on and completing tasks, or planning for and anticipating future events. Some exhibit learning difficulties and problems with language development.
- **Self-concept.** Traumatized children frequently suffer from disturbed body image, low self-esteem, shame, and guilt.

Long-Term Effects of Childhood Trauma

- In the absence of more positive coping strategies, children who have experienced trauma may engage in high-risk or destructive coping behaviors.
- These behaviors place them at risk for a range of serious mental and physical health problems, including:
 - Alcoholism
 - Drug abuse
 - Depression
 - Suicide attempts
 - Sexually transmitted diseases (due to high risk activity with multiple partners)
 - Heart disease, cancer, chronic lung disease, skeletal fractures, and liver disease

Childhood Trauma and PTSD

- Children who have experienced chronic or complex trauma frequently are diagnosed with PTSD.
- According to the American Psychiatric Association,¹ PTSD may be diagnosed in children who have:
 - Experienced, witnessed, or been confronted with one or more events that involved real or threatened death or serious injury to the physical integrity of themselves or others
 - Responded to these events with intense fear, helplessness, or horror, which may be expressed as disorganized or agitated behavior

Source: American Psychiatric Association. (2000).
DSM-IV-TR (4th ed.). Washington DC: APA.

20

Childhood Trauma and PTSD, cont'd

- Key symptoms of PTSD

- Reexperiencing the traumatic event (e.g. nightmares, intrusive memories)
- Intense psychological or physiological reactions to internal or external cues that symbolize or resemble some aspect of the original trauma
- Avoidance of thoughts, feelings, places, and people associated with the trauma
- Emotional numbing (e.g. detachment, estrangement, loss of interest in activities)
- Increased arousal (e.g. heightened startle response, sleep disorders, irritability)

Source: American Psychiatric Association. (2000).
DSM-IV-TR (4th ed.). Washington DC: APA.

Childhood Trauma and Other Diagnoses

- Other common diagnoses for children in the child welfare system include:
 - Reactive Attachment Disorder
 - Attention Deficit Hyperactivity Disorder
 - Oppositional Defiant Disorder
 - Bipolar Disorder
 - Conduct Disorder
- These diagnoses generally do not capture the full extent of the developmental impact of trauma.
- Many children with these diagnoses have a complex trauma history.

- Trauma can have serious consequences for the normal development of children's brains, brain chemistry, and nervous system.
- Trauma-induced alterations in biological stress systems can adversely effect brain development, cognitive and academic skills, and language acquisition.
- Traumatized children and adolescents display changes in the levels of stress hormones similar to those seen in combat veterans.
 - These changes may affect the way traumatized children and adolescents respond to future stress in their lives, and may also influence their long-term health.¹

1. Pynoos et al. (1997). *Ann N Y Acad Sci*;821:176-193

- In **early childhood**, trauma can be associated with reduced size of the cortex.
 - The cortex is responsible for many complex functions, including memory, attention, perceptual awareness, thinking, language, and consciousness.
- Trauma may affect “cross-talk” between the brain’s hemispheres, including parts of the brain governing emotions.
 - These changes may affect IQ, the ability to regulate emotions, and can lead to increased fearfulness and a reduced sense of safety and protection.

Trauma and the Brain, cont'd

- In **school-age children**, trauma undermines the development of brain regions that would normally help children:
 - Manage fears, anxieties, and aggression
 - Sustain attention for learning and problem solving
 - Control impulses and manage physical responses to danger, enabling the adolescent to consider and take protective actions
- As a result, children may exhibit:
 - Sleep disturbances
 - New difficulties with learning
 - Difficulties in controlling startle reactions
 - Behavior that shifts between overly fearful and overly aggressive

Trauma and the Brain, cont'd

- In **adolescents**, trauma can interfere with development of the prefrontal cortex, the region responsible for:
 - Consideration of the consequences of behavior
 - Realistic appraisal of danger and safety
 - Ability to govern behavior and meet longer-term goals
- As a result, adolescents who have experienced trauma are at increased risk for:
 - Reckless and risk-taking behavior
 - Underachievement and school failure
 - Poor choices
 - Aggressive or delinquent activity

Source: American Bar Association. (January 2004). Adolescence, Brain Development and Legal Culpability.

Available at: <http://www.abanet.org/crimjust/juvisus/Adolescence.pdf>

26

The Influence of Culture on Trauma

- Social and cultural realities strongly influence children's risk for—and experience of—trauma.
- Children and adolescents from minority backgrounds are at increased risk for trauma exposure and subsequent development of PTSD.
- In addition, children's, families' and communities' responses to trauma vary by group.

What Can a Provider Do?

- Understand that social and cultural realities can influence children's risk, experience, and description of trauma.
- Recognize that strong cultural identity can also contribute to resilience of children, their families, and their communities.
- Ensure that you become culturally competent and regularly participate in supervision.

What Can a Provider Do?, cont'd

- Consider how your own knowledge, experience, and cultural frame may influence your perceptions of traumatic experiences, their impact, and your choices of intervention strategies.
- Utilize resources the family trusts to supplement available services (e.g. bringing in a priest).

The Influence of Developmental Stage

- Child traumatic stress reactions vary by developmental stage.
- Children who have been exposed to trauma expend a great deal of energy responding to, coping with, and coming to terms with the event.
- This may reduce children's capacity to explore the environment and to master age-appropriate developmental tasks.
- The longer traumatic stress goes untreated, the farther children tend to stray from appropriate developmental pathways.

The Influence of Developmental Stage: Young Children

- **Young children** who have experienced trauma may:
 - Become passive, quiet, and easily alarmed
 - Become fearful, especially regarding separations and new situations
 - Experience confusion about assessing threat and finding protection, especially in cases where a parent or caretaker is the aggressor
 - Regress to recent behaviors (e.g., baby talk, bed-wetting, crying)
 - Experience strong startle reactions, night terrors, or aggressive outbursts

The Influence of Developmental Stage: School-Age Children

- **School-age children** with a history of trauma may:
 - Experience unwanted and intrusive thoughts and images
 - Become preoccupied with frightening moments from the traumatic experience
 - Replay the traumatic event in their minds in order to figure out what could have been prevented or how it could have been different
 - Develop intense, specific new fears linking back to the original danger

The Influence of Developmental Stage: School-Age Children, cont'd

- **School-age children** may also:
 - Alternate between shy/withdrawn behavior and unusually aggressive behavior
 - Become so fearful of recurrence that they avoid previously enjoyable activities
 - Have thoughts of revenge
 - Experience sleep disturbances that may interfere with daytime concentration and attention

The Influence of Developmental Stage: Adolescents

- In response to trauma, **adolescents** may feel:
 - That they are weak, strange, childish, or “going crazy”
 - Embarrassed by their bouts of fear or exaggerated physical responses
 - That they are unique and alone in their pain and suffering
 - Anxiety and depression
 - Intense anger
 - Low self-esteem and helplessness

The Influence of Developmental Stage: Adolescents, cont'd

- These trauma reactions may in turn lead to:
 - Aggressive or disruptive behavior
 - Sleep disturbances masked by late-night studying, television watching, or partying
 - Drug and alcohol use as a coping mechanism to deal with stress
 - Over- or under-estimation of danger
 - Expectations of maltreatment or abandonment
 - Difficulties with trust
 - Increased risk of revictimization, especially if the adolescent has lived with chronic or complex trauma

The Influence of Developmental Stage: Adolescents, Trauma, & Substance Abuse

- Adolescents who have experienced trauma may use alcohol or drugs in an attempt to avoid overwhelming emotional and physical responses. In these teens:
 - Reminders of past trauma may elicit cravings for drugs or alcohol.
 - Substance abuse further impairs their ability to cope with distressing and traumatic events.
 - Substance abuse increases the risk of engaging in risky activities that could lead to additional trauma.
- Child welfare workers must address the links between trauma and substance abuse and consider referrals for relevant treatment(s).

The Influence of Developmental Stage: Specific Adolescent Groups

- **Homeless youth** are at greater risk for experiencing trauma than other adolescents.
 - Many have run away to escape recurrent physical, sexual, and/or emotional abuse
 - Female homeless teens are particularly at risk for sexual trauma
- **Special needs adolescents** are 2 to 10 times more likely to be abused than their typically developing counterparts.
- **Lesbian, gay, bisexual, transgender or questioning (LGBTQ) adolescents** contend with violence directed at them in response to suspicion about or declaration of their sexual orientation and gender identity

What Can a Provider Do?

- Recognize that exposure to trauma is the rule, not the exception, among children in the child welfare system.
- Recognize the signs and symptoms of child traumatic stress and how they vary in different age groups.
- Recognize that children’s “bad” behavior is sometimes an adaptation to trauma.
- Understand the impact of trauma on different developmental domains.



What Can a Provider Do? cont'd


- Understand the cumulative effect of trauma.
- Gather and document psychosocial information regarding all traumas in the child's life to help their child welfare workers make better-informed decisions.
- Assist parents and caregivers who have secondary adversities and traumatic experiences of their own.
- Make a special effort to integrate cultural practices and culturally responsive interventions.
- Identify and build on foster parent and caregiver protective factors.

What Can a Provider Do?, cont'd

- Recognize that child welfare system interventions have the potential to either exacerbate or decrease the impact of previous traumas.
- Lessen the risk of system-induced secondary trauma by serving as a protective and stress-reducing buffer for children:
 - Develop *trust* with children through listening, frequent contacts, and honesty in order to mitigate previous traumatic stress.
 - Avoid repeated interviews, especially about experiences of sexual abuse.
 - Avoid making professional promises that, if unfulfilled, are likely to increase traumatization.

Child Welfare Trauma Training :

The Impact of Trauma on Children's Behavior, Development, and Relationships



Artwork courtesy of the International Child Art Foundation (www.icaf.org)

Maximize the child's sense of safety

- Traumatic stress overwhelms a child's sense of safety and can lead to a variety of survival strategies for coping.
- Safety implies both *physical* safety and *psychological* safety.
- A sense of safety is critical for functioning as well as physical and emotional growth.
- While inquiring about emotionally painful and difficult experiences and symptoms, workers must ensure that children are provided a psychologically safe setting.

Maximizing Safety: Understanding Children's Responses

- Children who have experienced trauma often exhibit extremely challenging behaviors and reactions.
- When we label these behaviors as “good” or “bad,” we forget that children’s behavior is reflective of their experience.
- Many of the most challenging behaviors are strategies that in the past may have helped the child survive in the presence of abusive or neglectful caregivers.

Assist children in reducing overwhelming emotion

- Trauma can elicit such intense fear, anger, shame, and helplessness that the child feels overwhelmed.
- Overwhelming emotion may delay the development of age-appropriate self-regulation.
- Emotions experienced prior to language development maybe be very real for the child but difficult to express or communicate verbally.
- Trauma may be “stored” in the body in the form of physical tension or health complaints.

Reduce Overwhelming Emotion: Understanding Trauma Reminders

- When faced with people, situations, places, or things that remind them of traumatic events, children may experience intense and disturbing feelings tied to the original trauma.
 - These “**trauma reminders**” can lead to behaviors that seem out of place, but were appropriate—and perhaps even helpful—at the time of the original traumatic event.
- Children who have experienced trauma may face so many trauma reminders in the course of an ordinary day that the whole world seems dangerous and no adult seems deserving of trust.

Reduce Overwhelming Emotion: Understanding Children's Responses

- When placed in a new, presumably “safe” setting, traumatized children may exhibit behaviors (e.g., aggression, sexualized behaviors) that evoke in their new caregivers some of the same reactions they experienced with other adults (e.g., anger, threats, violence).
- Just as traumatized children's sense of themselves and others is often negative and hopeless, these “**reenactment behaviors**” can cause the new adults in their lives to feel negative and hopeless about the child.

Reduce Overwhelming Emotion: Understanding Children's Responses, cont'd

- Children who engage in reenactments are not consciously choosing to repeat painful relationships. The behavior patterns have become ingrained over time because they:
 - Are familiar and helped the child survive in other relationships
 - “Prove” the child’s negative beliefs and expectations (a predictable world, even if negative, may feel safer than an unpredictable one)
 - Help the child vent frustration, anger, and anxiety
 - Give the child a sense of mastery over the old traumas

47

Reduce Overwhelming Emotion: Understanding Children's Responses, cont'd

- Traumatized children may also exhibit:
 - **Over-controlled behavior** in an unconscious attempt to counteract feelings of helplessness and impotence
 - May manifest as difficulty transitioning and changing routines, rigid behavioral patterns, repetitive behaviors, etc.
 - **Under-controlled behavior** due to cognitive delays or deficits in planning, organizing, delaying gratification, and exerting control over behavior
 - May manifest as impulsivity, disorganization, aggression, or other acting-out behaviors

Reduce Overwhelming Emotion: Understanding Children's Responses, cont'd

- Traumatized children's maladaptive coping strategies can lead to behaviors that undermine healthy relationships and may disrupt foster placements, including:
 - Sleeping, eating, elimination problems
 - High activity level, irritability, acting out
 - Emotional detachment, unresponsiveness, distance, or numbness
 - Hypervigilance or feeling that danger is present, even when it isn't
 - Increased mental health issues (e.g. depression, anxiety)
 - An unexpected and exaggerated response when told "no"

49

Reduce Overwhelming Emotion: What Providers Can Do

- When appropriate, conduct a trauma-focused mental health assessment to identify services and interventions appropriate to the child's needs.
- Share the child's traumatic experiences and anticipated responses with foster placement as appropriate.
- Encourage foster parents to provide information if/when new revelations of past traumas emerge.
- Empower caregivers about their role of calming and reassuring children.

Reduce Overwhelming Emotion: What Providers Can Do, cont'd

- Educate caregivers about the reasons for, and techniques to manage, children's emotional outbursts.
- Recommend parenting skills training to strengthen caregivers' ability to handle children's emotions.
- Work with the child to identify and label troubling emotions and stress that the emotions are normal and understandable.

Help children make new meaning of their trauma history and current experiences

- Trauma can lead to serious disruptions in a child's sense of safety, personal responsibility, and identity.
- Distorted connections between thoughts, feelings, and behaviors can disrupt encoding and processing of memory.
- Difficulties in communicating about the event may undermine child's confidence and social support.
- Providers can assist traumatized children in developing a coherent understanding of their traumatic experiences.

Make New Meaning of Trauma History: What Providers Can Do

- When possible, gather a complete trauma history from child welfare workers, caregivers and/or child.
- As appropriate, provide the child with information about events that led to child welfare involvement in order to help the child correct distortions and reduce self-blame.
- Listen to and acknowledge the child's traumatic experience(s).

Make New Meaning of Trauma History: What Providers Can Do, cont'd

- Support the child in the development of a Life Book (i.e., a book of stories and memories about the child's life).
- Receive training/educate yourself on evidence-based trauma-focused therapies and be open to including these new techniques into your current treatment modality.
- Try to include current caregivers in treatment and educate them about the impact of trauma on child behaviors and behavior management.

Child Welfare Trauma Training:

Assessment of a Child's Trauma Experiences



Artwork courtesy of the International Child Art Foundation (www.icaf.org)

Address the impact of trauma:

- Trauma affects many aspects of the child's life and can lead to secondary problems (e.g., difficulties in school and relationships, or health-related problems).
- These “secondary adversities” may mask symptoms of the underlying traumatic stress and interfere with a child's recovery from the initial trauma.
- Secondary adversities can also lead to changes in the family system and must be addressed prior to or along with trauma-focused interventions.

Coordinate services with other agencies:

- Traumatized children and their families are often involved with multiple service systems.
- Cross-system collaboration enables all helping professionals to see the child as a whole person, thus preventing potentially competing priorities and messages.
- Providers should try to develop common protocols and frameworks for documenting trauma history, exchanging information, coordinating assessments, and planning and delivering care.

Utilize comprehensive assessment:

- Trauma-specific standardized assessments can identify potential risk behaviors (i.e. danger to self, danger to others) and help determine interventions that will reduce risk.
- Thorough assessment can identify a child's reactions and how his or her behaviors are connected to the traumatic experience.
- Assessment results provide valuable information for developing treatment goals with measurable objectives designed to reduce the negative effects of trauma.
- Assessment results also can be used to determine the need for referral to trauma-specific mental health care or more detailed trauma assessment.

The Importance of Trauma Assessment

- Not all children who have experienced trauma need trauma-specific intervention.
- Some children have amazing natural resilience and are able to use their natural support systems to integrate their traumatic experience.
- Ideally, children should be in a stable placement when receiving trauma-informed treatment. However, children should *always* be referred for necessary treatment regardless of their placement status.

The Importance of Trauma Assessment, cont'd

- Unfortunately, many children in the child welfare system lack natural support systems and need the help of trauma-informed care. Some may meet the clinical criteria for a diagnosis of PTSD.
- Many children who do not meet the full criteria for PTSD still suffer significant posttraumatic symptoms that can have a dramatic adverse impact on behavior, judgment, educational performance, and ability to connect with caregivers.
- These children need a comprehensive trauma assessment to determine which intervention will be most beneficial.

The Importance of Trauma Assessment, cont'd

- Trauma assessment typically involves conducting a thorough trauma history.
 - Identify all forms of traumatic events experienced directly or witnessed by the child to determine the best type of treatment for that specific child.
- Supplement trauma history with trauma-specific standardized clinical measures to assist in identifying the types and severity of symptoms the child is experiencing.

What Does Trauma-Informed Assessment and Treatment Look Like?

- There are evidence-supported interventions that are appropriate for many children and that share many core components of trauma-informed treatments.
- Unfortunately, many therapists who treat traumatized children lack any specialized knowledge or training on trauma and its treatment.
- “Trauma” and “Trauma Treatment” are becoming important words in the field of child welfare and workers and agencies are seeking therapists who have had any training and/or skills in this area.

Examples of Trauma Assessment Measures

- UCLA PTSD Index for DSM-IV
- Trauma Symptom Checklist for Children (TSCC)
- Trauma Symptom Checklist for Young Children (TSCYC)
- Child Sexual Behavior Inventory

Core Components of Trauma-Informed, Evidence-Based Treatment

- Building a strong therapeutic relationship
- Psychoeducation about normal responses to trauma
- Parent support, conjoint therapy, or parent training
- Emotional expression and regulation skills
- Anxiety management and relaxation skills
- Cognitive processing or reframing

Core Components of Trauma-Informed, Evidence-Based Treatment, cont'd

- Construction of a coherent trauma narrative
- Strategies that allow exposure to traumatic memories and feelings in tolerable doses so that they can be mastered and integrated into the child's experience
- Personal safety training and other important empowerment activities
- Resilience and closure

Questions to Ask Yourself and/or other Providers who provide Trauma Treatment

- Do you provide trauma-specific or trauma-informed therapy? If so, how do you determine if the child needs a trauma-specific therapy?
- How familiar are you with evidence-based treatment models designed and tested for treatment of child trauma-related symptoms?
- How do you approach therapy with traumatized children and their families (regardless of whether they indicate or request trauma-informed treatment)?
- Describe a typical course of therapy (e.g., can you describe the core components of your treatment approach?).

Examples of Evidence-Based Treatments

- Trauma-Focused Cognitive Behavioral Therapy (TF-CBT)
- Parent-Child Interaction Therapy (PCIT)
- Abuse-Focused Cognitive Behavioral Therapy (AF-CBT)
- Child-Parent Psychotherapy (CPP)

There are many different evidence-based trauma-focused treatments. A trauma-informed mental health professional should be able to determine which treatment is most appropriate for a given case.

Trauma-Focused Cognitive Behavioral Therapy (TF-CBT)

- Originally developed to treat child sexual abuse
- An empirically supported intervention based on learning and cognitive theories
- Designed to reduce children's negative emotional and behavioral responses, and to correct maladaptive beliefs and attributions related to the abusive experiences
- Aims to provide support and skills to help non-offending parents cope effectively with their own emotional distress and to respond optimally to their abused children

Cohen, et al. (2006). *Treating Trauma and Traumatic Grief in Children and Adolescents*. New York: Guilford Press.



Core Components of TF-CBT

- Stress inoculation techniques
 - Feeling identification
 - Relaxation, thought stopping, cognitive coping
- Cognitive processing (part 1)
 - The cognitive triangle
- Creating a trauma narrative
- Cognitive processing (part 2)
 - Processing the trauma experience
 - Joint family sessions
 - Psychoeducation

Parent-Child Interaction Therapy (PCIT)

- Works with the caregiver and child together
- Designed to treat children aged 2–8 years who are exhibiting disruptive behaviors
- Use of coaching: caregiver wears hidden earpiece and is prompted by therapist behind a one-way mirror
- Average of 14–20 weekly sessions focused on relationship enhancement and behavior management
- Combines elements from family systems, operant, social learning, and traditional play therapies, as well as early child development theory

Child-Parent Psychotherapy (CPP)

- Works with the caregiver and child together.
- Designed to treat children aged 0–6 years.
- Based in attachment theory but also integrates psychodynamic, developmental, trauma, social learning, and cognitive behavioral theories.
- Focus is on safety, affect regulation, improving the child caregiver relationship, normalization of trauma related response, joint construction of a trauma narrative, with the goal of returning the child to a normal developmental trajectory
- Studies have shown significant reduction in behavior problems and traumatic stress symptoms and in improved representation of self and caregiver. With babies/toddlers, studies show an increase in secure attachment after treatment.

Lieberman, AF, and Van Horn, P. (2005). "Don't hit my mommy!": A manual for child-parent psychotherapy with young witnesses of family violence.


Washington, DC: Zero to Three Press.



What Providers Can Do:

- To obtain training/mentoring, find other providers or agencies who are knowledgeable about trauma assessment and evidence-based treatments; join local professional groups.
- Gather a full picture of a child's experiences and trauma history.
- Identify immediate needs and concerns in order to prioritize interventions for specific individuals.
- Conduct regular, ongoing assessments (e.g., every three months) regarding the child's progress and symptoms.

Child Welfare Trauma Training : Managing Professional and Personal Stress



Artwork courtesy of the International Child Art Foundation (www.icaf.org)

- Therapists/counselors may empathize with their clients' experiences; feelings of helplessness, anger, and fear are common.
- Therapists/counselors who are parents—or who have their own histories of childhood trauma—may be at particular risk for experiencing such reactions.
- Regular supervision and/or therapy is essential to maintain objectivity and for managing stress.

Impact of Working with Victims of Trauma

- Trauma experienced while working in the role of helper has been described as:
 - Compassion fatigue
 - Countertransference
 - **Secondary traumatic stress (STS)**
 - Vicarious traumatization
- Unlike other forms of job “burnout,” STS is precipitated not by work load and institutional stress but by exposure to clients’ trauma.
- STS can disrupt provider’s lives, feelings, personal relationships, and overall view of the world.

Managing Stress: What Providers Can Do

- Request and expect regular supervision and supportive consultation.
- Utilize peer support.
- Consider therapy for unresolved trauma, which the work may be activating.
- Practice stress management through meditation, prayer, conscious relaxation, deep breathing, and exercise.
- Develop a written plan focused on maintaining work–life balance.



Thank You!

